

UM-TH-93-20

IP-ASTP-29-93

August 25, 1993

Exact α_s Calculation of $b \rightarrow s + \gamma$

$$b \rightarrow s + g$$

K. Adel^a and York-Peng Yao^{a,b}^a *Department of Physics, Randall Laboratory, University of Michigan,**Ann Arbor, Michigan 48109 – 1120, U.S.A.*^b *Institute of Physics, Academia Sinica, Taipei,**Taiwan 11529, Republic of China.**Abstract:*

We present an exact α_s calculation of the Wilson coefficients associated with the dipole moment operators. We also give an estimate of the branching ratio for $b \rightarrow s\gamma$. We find that higher dimensional effects are under control within 9% for $BR(b \rightarrow s\gamma) = (4.3 \pm 0.37) \times 10^{-4}$.

PAC: 11.10Jj, 12.38.Bx, 13.40.Hq

In an article¹ by us recently, we presented a leading logarithmic analysis of the heavy particle effects on the process $b \rightarrow s\gamma$, which incorporates a complete operator mixing. Among the results, we find that our mixing matrix differs in some elements from the work by Misiak². Had we subscribed to the prescription of further mixing with evanescent operators³, we would have obtained extra contributions (in unit of $g_s^2/8\pi^4$)

$$\begin{aligned}\Delta\gamma_{O_{67} O_{51}} &= 1/3, & \Delta\gamma_{O_{67} O_{52}} &= -8/3, \\ \Delta\gamma_{O_{68} O_{51}} &= -1/2, & \Delta\gamma_{O_{68} O_{52}} &= -4.\end{aligned}\tag{1}$$

These would have made the mixing by Ciuchini et. al.⁴ coincide with ours. (We thank Misiak for correspondence on this.) For earlier work, see^{5,6}.

More importantly, by choosing two different limits for extrapolation: $m_t = m_w$ and $m_t \gg m_w$, we estimated the effects due to higher orders in m_w^2/m_t^2 to be about 20%. In view of recent interests in the experimental branching ratio⁷ for $B_s \rightarrow K^* + \gamma$ and an impending value for the inclusive rate $B_s \rightarrow X_s + \gamma$, together with the sensitivity of these processes as a short distance probe, an uncertainty in short distance analysis of 20% is hardly satisfactory or even acceptable. The purpose of this publication is to provide some remedy.

One can trace the uncertainty to the fact that $m_w^2/m_t^2 \approx 25\%$ for $m_t \approx 150$ Gev. It strongly suggests that one should calculate the order α_s diagrams exactly, wherever there are internal top and/or W-boson propagators. This is what we have done.

In all two-loop diagrams (and their attendant counter terms) which contribute to this calculation, we keep all orders in $x \equiv m_t^2/m_w^2$ but discard terms proportional to $O(m_b^2/m_{t,w}^2)$ or $O(m_s^2/m_{t,w}^2)$ (we have factored out the Fermi weak coupling constant). The amount of algebra is highly non-trivial and is aided by Schoonschip.⁸

Given a diagram in which there are top and/or $W - \phi$ internal lines, there are various sequences of operations one can follow to isolate its dependence on the heavy masses⁹. In any case, the underlying method is based on partitioning of the diagrams into heavy parts and operator inserted matrix elements. For the present situation, we treat the top and the $W - \phi$ as being correspondingly heavy, relative to other masses and external momenta. In the order we are working with, a heavy part always contains both the top and the $W - \phi$ internal lines. Vertices made of light particles and momenta acting on them and Wilson coefficients which contain all dependence of heavy masses are organized within this formalism. In this way, we obtain the effective Lagrangian

$$L_{eff} = \sum_i C_i O_i, \quad (2)$$

where C_i are the Wilson coefficients and O_i are a set of local operators, made of light fields.

Of particular interest in the present order $\alpha_w \alpha_s$ calculation are the coefficients $C_{O_{51}}$ and $C_{O_{52}}$, with the accompanying operators

$$O_{51} = ig_s \bar{s} G_{\mu\nu} \sigma_{\mu\nu} (m_s P_L + m_b P_R) b / 2, \quad (3)$$

and $O_{52} = O_{51} (g_s G_{\mu\nu} \rightarrow (-1/3) e F_{\mu\nu})$.

Before we go on further, let us define

$$\bar{C}_{O_{5i}} \equiv 16\pi^2 C_{O_{5i}} / G_t \quad (i = 1, 2), \quad \text{and} \quad G_t = 2\sqrt{2} G_F V_{ts}^* V_{tb}, \quad (4)$$

where G_F is the Fermi weak coupling constant and V 's are the Cabbibo-Kobayashi-

Maskawa matrix elements. Then the exact α_s results are (in the R^* scheme to be defined)

$$\begin{aligned}
\frac{\pi}{4\alpha_s} \bar{C}_{O_{51}}^{(1)} = & -\frac{4}{(x-1)^4} - \frac{5293}{576} \frac{1}{(x-1)^3} - \frac{21989}{3456} \frac{1}{(x-1)^2} - \frac{1817}{1728} \frac{1}{(x-1)} + \frac{247}{10368} \\
& + \log\left(\frac{m_w^2}{\mu^2}\right) \left[-\frac{7}{16} \frac{1}{(x-1)^3} - \frac{21}{32} \frac{1}{(x-1)^2} - \frac{7}{48} \frac{1}{(x-1)} - \frac{35}{288} \right] \\
& + \log\left(\frac{m_w^2}{\mu^2}\right) \log x \left[\frac{7}{16} \frac{1}{(x-1)^4} + \frac{7}{8} \frac{1}{(x-1)^3} + \frac{7}{16} \frac{1}{(x-1)^2} \right] \\
& + \log x \left[\frac{4}{(x-1)^5} + \frac{5509}{576} \frac{1}{(x-1)^4} + \frac{2893}{432} \frac{1}{(x-1)^3} + \frac{163}{192} \frac{1}{(x-1)^2} - \frac{11}{48} \frac{1}{(x-1)} - \frac{35}{288} \right] \\
& + \log^2 x \left[+\frac{7}{16} \frac{1}{(x-1)^4} + \frac{7}{8} \frac{1}{(x-1)^3} + \frac{7}{16} \frac{1}{(x-1)^2} \right] \\
& + \text{Sp}\left(1 - \frac{1}{x}\right) \left[\frac{13}{8} \frac{1}{(x-1)^4} + \frac{187}{48} \frac{1}{(x-1)^3} + \frac{137}{48} \frac{1}{(x-1)^2} + \frac{1}{2} \frac{1}{(x-1)} - \frac{1}{12} \right] \\
& + \left(\frac{91}{2592} + \frac{1}{54} \log\left(\frac{m_w^2}{\mu^2}\right) \right), \\
\\
\frac{\pi}{4\alpha_s} \bar{C}_{O_{52}}^{(1)} = & -\frac{4}{(x-1)^4} - \frac{335}{18} \frac{1}{(x-1)^3} - \frac{11959}{432} \frac{1}{(x-1)^2} - \frac{6347}{432} \frac{1}{(x-1)} - \frac{47}{648} \\
& + \log\left(\frac{m_w^2}{\mu^2}\right) \left[-\frac{1}{(x-1)^3} - \frac{3}{(x-1)^2} - \frac{31}{12} \frac{1}{(x-1)} + \frac{17}{36} \right] \\
& + \log\left(\frac{m_w^2}{\mu^2}\right) \log x \left[\frac{1}{(x-1)^4} + \frac{7}{2} \frac{1}{(x-1)^3} + \frac{4}{(x-1)^2} + \frac{3}{2} \frac{1}{(x-1)} \right] \\
& + \log x \left[\frac{4}{(x-1)^5} + \frac{643}{36} \frac{1}{(x-1)^4} + \frac{1337}{54} \frac{1}{(x-1)^3} + \frac{257}{24} \frac{1}{(x-1)^2} - \frac{19}{24} \frac{1}{(x-1)} + \frac{17}{36} \right] \\
& + \log^2 x \left[\frac{1}{(x-1)^4} + \frac{7}{2} \frac{1}{(x-1)^3} + \frac{4}{(x-1)^2} + \frac{3}{2} \frac{1}{(x-1)} \right] \\
& + \text{Sp}\left(1 - \frac{1}{x}\right) \left[\frac{11}{4} \frac{1}{(x-1)^4} + \frac{139}{12} \frac{1}{(x-1)^3} + \frac{191}{12} \frac{1}{(x-1)^2} + \frac{31}{4} \frac{1}{(x-1)} + \frac{2}{3} \right] \\
& + \left(-\frac{713}{648} + \frac{1}{54} \log\left(\frac{m_w^2}{\mu^2}\right) \right),
\end{aligned} \tag{5}$$

where the last terms in big parenthesis in each of the two equations above are the contributions due to u and c quarks. Sp stands for Spence function. Eq.(5) are to be contrasted with the asymptotic results ($m_t \gg m_w$)

$$\bar{C}_{O_{51}}^{(1)}(asym)\pi/4\alpha = \frac{611}{10368} - \frac{\pi^2}{72} - \frac{35}{288} \log\left(\frac{m_t^2}{\mu^2}\right) + \frac{1}{54} \log\left(\frac{m_w^2}{\mu^2}\right),$$

and

$$\bar{C}_{O_{52}}^{(1)}(asym)\pi/4\alpha = -\frac{95}{81} + \frac{\pi^2}{9} + \frac{17}{36} \log\left(\frac{m_t^2}{\mu^2}\right) + \frac{1}{54} \log\left(\frac{m_w^2}{\mu^2}\right), \quad (6)$$

obtained by discarding all $O(m_w^2/m_t^2)$. One can use Eq.(5) to check some of the mixing matrix elements of the renormalization group equations (RGE). Unfortunately, $\gamma_{O_{67} O_{51}}$, $\gamma_{O_{67} O_{52}}$, $\gamma_{O_{68} O_{51}}$, and $\gamma_{O_{68} O_{52}}$ do not enter to this order. Their determination has to come from three loop diagrams. In our opinion, direct Feynman diagram computation of Green's functions for processes to extract mixing matrix elements should be the definitive procedure.

We plot the exact α_s result Eq.(5) in Figure 1, together with the asymptotic result Eq.(6) for $\bar{C}_{O_{52}}^{(1)}\pi/4\alpha_s$. For $x=4$, the discrepancy is in fact about 50%. It is interesting to note that the exact $\bar{C}_{O_{52}}^{(1)}$ is quite flat between $x=1$ to $x=10$. The dotted line is the α_s^0 exact result, also known as the Inami-Lim functions¹⁰

$$\begin{aligned} \bar{C}_{O_{51}}^{(0)} &= \frac{-2x - 3x^2 + 6x^3 - x^4 - 6x^2 \log x}{4(1-x)^4}, \\ \bar{C}_{O_{52}}^{(0)} &= \frac{7x - 12x^2 - 3x^3 + 8x^4 + (12x^2 - 18x^3) \log x}{4(1-x)^4}. \end{aligned} \quad (7)$$

We see that below $x=6.9$, second order QCD correction is bigger than the lowest order result. This has been known for the approximate results for some time and in fact is an impetus for looking into rare decays of this genre.

Please note that if one uses Eq.(7) as (a part of) the boundary conditions, then Eq.(5) are the α_s solution of RGE for all values of x , in so far as the overall coefficients to $\log \mu^2$ are concerned. We would like to stress that we are using the R^* scheme. In the $\overline{MS}$ scheme, the exact result to order α_s is obtained if one makes the replacement $m_t \rightarrow m_t (1 - \alpha_s/\pi \log(m_t^2/\mu^2))$ in Eq. (7); this replacement is just a finite renormalization to go from the R^* scheme to the $\overline{MS}$ scheme.

We shall make the assumption that after α_s corrections, it is safe to add the leading logarithmic terms to complete the leading QCD sum. In other words, we assume that the higher order QCD corrections can be obtained either in the limit $m_w^2/m_t^2 \ll 1$ or in the limit $m_w^2/m_t^2 = 1$. This assumption can be tested as in our previous publication. For $\bar{C}_{O_{52}}$, there entail two different extrapolations

$$\bar{C}_{O_{52}}^{(0)+(1)}(exact) + \bar{C}_{O_{52}}^{higher\ order}(m_t = m_w) \quad (8.a),$$

and

$$\bar{C}_{O_{52}}^{(0)+(1)}(exact) + \bar{C}_{O_{52}}^{higher\ order}(m_t \gg m_w), \quad (8.b)$$

where $\bar{C}_{O_{52}}^{higher\ order}(m_t = m_w)$ and $\bar{C}_{O_{52}}^{higher\ order}(m_t \gg m_w)$ are the remaining leading logarithmic sums with the boundary conditions set at $m_t = m_w$ and $m_t \gg m_w$, respectively.¹

For the physical process $b \rightarrow s + \gamma$, some four quark operators also contribute, resulting in an effective coupling²

$$C_{O_{52}}^{eff} = C_{O_{52}} + \frac{1}{8\pi^2} C_{O_{67}} + \frac{3}{8\pi^2} C_{O_{68}}. \quad (9)$$

Also, to remove the dependence on $|G_t|^2$, which is not accurately known experimentally, we normalize the $b \rightarrow s\gamma$ partial width to the well established semileptonic $b \rightarrow ce\bar{\nu}$ partial width, and use the following relation¹¹ $|V_{ts}^* V_{tb}| \simeq |V_{cb}|$.

This ratio is given as:

$$\frac{\Gamma(b \rightarrow s\gamma)}{\Gamma(b \rightarrow ce\bar{\nu})} \simeq \frac{\alpha_{QED}}{6\pi g(m_c/m_b)} \left(1 - \frac{2\alpha_s(m_b)}{3\pi} f(m_c/m_b)\right)^{-1} |\bar{C}_{O_{52}}^{eff}(m_b)|^2, \quad (10)$$

where $g(m_c/m_b) \simeq 0.45$ and $f(m_c/m_b) \simeq 2.4$ correspond to the phase space factor and the one-loop QCD corrections to the semileptonic decay, respectively.

In Figure 2, we have plotted $BR(b \rightarrow s\gamma)$ as a function of m_t . The solid and dashed curves are obtained with the aid of the interpolation equations in Eq.(8), together with:

$$BR(b \rightarrow s\gamma) = \frac{\Gamma(b \rightarrow s\gamma)}{\Gamma(b \rightarrow ce\bar{\nu})} BR(b \rightarrow ce\bar{\nu}), \quad (11)$$

$$BR(b \rightarrow ce\bar{\nu}) \simeq 0.108. \quad (12)$$

The vertical dotted line to guide the eyes intersects these curves at $m_t = 140 \text{ GeV}$ and gives respectively:

$$\begin{aligned} BR(b \rightarrow s\gamma) &= 4.66 \times 10^{-4}, \\ BR(b \rightarrow s\gamma) &= 3.93 \times 10^{-4}. \end{aligned} \quad (13)$$

From Eq. (13), we obtain the following mean value (this is our estimate),

$$BR(b \rightarrow s\gamma) = (4.3 \pm 0.37) \times 10^{-4}. \quad (14)$$

This is to be compared with an upper limit 5.4×10^{-4} given recently by the CLEO Collaboration.⁽⁷⁾

The uncertainty due to subleading logarithmic and higher dimensional effects is about 9%, which is a big improvement and more reliable over what we gave before, where m_t^2/m_w^2 effects at $\alpha_s^{(1)}$ were not treated.

We now give some technical details. We use the general linear covariant gauges $\frac{-1}{2\alpha}(\partial_\mu G_\mu)^2$ for the gluons. The complete cancellation of α for $C_{O_{51}}$ and $C_{O_{52}}$ is a stringent confirmation on the correctness of the algebra. The gauge fixing for W-fields is $-C^+C^-$ with $C^+ = -\partial_\mu W_\mu^+ + m_w \phi^+ + ieA_\mu W_\mu$. For oversubtractions and renormalization, we use the R^* -scheme. Thus, let Γ be a one light particle irreducible (1LPI) diagram which contains the heavy particles, and let γ represent a 1LPI graph or subgraph ($\gamma \subseteq \Gamma$) with external generic momentum p . We define

$$\begin{aligned} \tau_\gamma^\epsilon &= \text{pole part of } \gamma, \quad \epsilon = n - 4, \\ \tau_\gamma^{(m)} &= \gamma(p=0) + p \frac{\partial}{\partial p} \gamma(p=0) + \cdots + \frac{1}{m!} p^m \frac{\partial^m}{\partial p^m} \gamma(p=0). \end{aligned} \quad (11)$$

The R^* renormalization procedure is defined as

$$R^*(\gamma_{heavy}) = (1 - \tau_\gamma^{(m)})\gamma_{heavy}, \quad R^*(\gamma_{light}) = (1 - \tau_\gamma^\epsilon)\gamma_{light}, \quad (12)$$

where m is so chosen that the neglected terms are genuinely of $O(1/m_{heavy}^2)$. It is important to repeat that the results Eqs.(5,6) are given in the renormalized top mass under the R^* -scheme, where $\beta_{m_t} = 0$.

Except for trivial factorizable cases, all two loop integrals we need are related to¹²

$$\begin{aligned}
I_{2,1,1}(m_1^2, m_2^2, m_3^2) &= \int d^n k d^n q \frac{1}{(k^2 + m_1^2)^2 ((k+q)^2 + m_2^2) (q^2 + m_3^2)} \\
&= \pi^4 \left[\frac{-2}{(n-4)^2} + \frac{1}{n-4} (1 - 2\gamma_E - 2\log(\pi m_1^2)) - \frac{1}{2} - \frac{1}{12}\pi^2 \right] \\
&\quad + \pi^4 \left[\gamma_E - \gamma_E^2 + (1 - 2\gamma_E) \log(\pi m_1^2) - \log^2(\pi m_1^2) + f(a = \frac{m_2^2}{m_1^2}, b = \frac{m_3^2}{m_1^2}) \right],
\end{aligned} \tag{13}$$

where

$$\begin{aligned}
f(a, b) &= \frac{1}{2} \log a \log b + \frac{a+b-1}{\sqrt{}} [\text{Sp}(\frac{-y_2}{x_1}) + \text{Sp}(\frac{-x_2}{y_1}) + \frac{1}{4} \log^2 \frac{x_2}{y_1} \\
&\quad + \frac{1}{4} \log^2 \frac{y_2}{x_1} + \frac{1}{4} \log^2 \frac{x_1}{y_1} - \frac{1}{4} \log^2 \frac{x_2}{y_2} + \frac{\pi^2}{6}],
\end{aligned} \tag{14}$$

$$x_{1,2} = \frac{1}{2}(1 - a + b \pm \sqrt{}), \quad y_{1,2} = \frac{1}{2}(1 + a - b \pm \sqrt{}),$$

and

$$\sqrt{} = \sqrt{(1-a+b)^2 - 4b}. \tag{15}$$

Assuming $a, b \gg 1$, (i. e. we take $m_2 = m_t$, $m_3 = m_w \gg m_1 = m_{b,s}$), we expand $f(a, b)$ in series of $m_{b,s}^2$, which must be retained to proper orders in intermediate steps.

Details of this work are to be published¹.

This work has been partially supported by the U. S. Department of Energy. Y.-P. Y. would like to thank members of the Particle Theory Group at the Institute of Physics, Academia Sinica, for hospitality.

References:

- [1] K. Adel and York-Peng Yao, Modern Physics Letters A **8**, 1679 (1993);
also under preparation.
- [2] M. Misiak, Phys. Lett. **269 B**, 161 (1991); Nucl. Phys. **B393**, 23 (1993).
- [3] A. Buras and P. Weisz, Nucl. Phys. **B333**, 66 (1990).
- [4] M. Ciuchini, E. Franco, G. Martinelli, L. Reina, L. Silvestrini, LPTENS 93/28,
ROME 93/958, ULB-TH 93/09 (unpublished).
- [5] B. Grinstein, R. Springer and M. B. Wise, Phys. Lett. **202 B**, 138 (1988);
P. Cho and B. Grinstein, Nucl. Phys. **B365**, 138 (1991);
See also references listed in these articles.
- [6] R. Griganis, P. J. O'Donnell, M. Sutherland and H. Navalet, Phys. Lett. **213 B**,
355 (1988); R. Griganis, P. J. O'Donnell and M. Sutherland, Phys. Lett. **237 B**,
252 (1990); P. J. O'Donnell and H. K. K. Tung, Phys. Rev. **D 45**, 252 (1990);
G. Cella, G. Curci, G. Ricciardi and A. Viceré, Phys. Lett. **248 B**, 181 (1990).
- [7] E. Thorndike, CLEO Collaboration, talk given at the *1993 Meeting of the American
Physical Society, Washington, D. C. April, 1993*.
- [8] Schoonschip program by M. Veltman, unpublished.
- [9] W. Zimmermann, in *Lectures in Elementary Particles and Quantum Field
Theory*, edited by S. Deser *et al.* (MIT Press, Cambridge, Mass., 1971),
Vol. I, p.397; Y. Kazama and Y.-P. Yao, Phys. Rev. **D 25**, 1605 (1982).
- [10] T. Inami and C. S. Lim, Progr. Theor. Phys. **65** 297 (1981).
- [11] N. Cabibbo and L. Maiani, Phys. Lett. **79 B**, 109 (1978);
B. A. Campbell and P. J. O'Donnell, Phys. Rev. **D 25**, 1989 (1982).
- [12] J. van der Bij and M. Veltman, Nucl. Phys. **B231**, 205 (1984).

FIGURE CAPTION:

Figure 1: $\bar{C}_{O_{52}}(m_b)$ dependence on m_t to order α_s with $m_b = 4.8 \text{ GeV}$, $m_w = 81 \text{ GeV}$, $\alpha_s(m_b) = 0.19$. See text for explanation of various curves.

Figure 2: Branching ratio for $b \rightarrow s\gamma$ as a function of m_t with QCD corrections. The solid line represents the interpolation given by Eq. (8.a); the dashed line represents the interpolation given by Eq. (8.b); the dotted line represents the values of the branching ratio for $m_t = 140 \text{ GeV}$. We used $m_b = 4.8 \text{ GeV}$, $m_w = 81 \text{ GeV}$, $\alpha_s(m_b) = 0.19$.